Physics 681: Solar Physics and Instrumentation – Lecture 12

Carsten Denker

NJIT Physics Department Center for Solar–Terrestrial Research

Oscillations

Observations

- Five-Minute Oscillation
- The Spectrum of Solar Oscillations
- Low-Degree p Modes
- Line Width and Asymmetry
- Linear Adiabatic Oscillations of a Non-Rotating Sun
 - Basic Equations
 - Spherical Harmonic Representation
 - The Cowling Approximation
 - Local Treatment
 - Boundary Conditions
 - Asymptotic Results
- Helioseismology
- Excitation and Damping

October 13, 2005

Center for Solar-Terrestrial Research



- Acoustic waves
- Pressure gradient is the restoring force "p modes"
- Probe the interior of the Sun
- Gravity can also provide a restoring force
- However, g modes have not yet been confirmed
- Leighton et al. discovered the fiveminute oscillations in 1960 in dopplergrams
- Spectroscopy ⇒ vertical velocities from 0.5 to 1.0 km/s
- Periodic variations of the intensity (few percent)
- Largest amplitudes (2-5 mHz or 3-8 min)
- Areas of 30,000 km oscillate in phase
- Evanescent waves

October 13, 2005



Spectrum of Solar Oscillations

Temporal and spatial frequencies

 $\Delta \omega = 2\pi / T \le \omega < \pi / \Delta t$ $\underline{\Delta k_x} = 2\pi / L_x \le \omega < \pi / \Delta x$

□ Fourier transform of a vertical velocity signal $v(x, y, t) = \int f(k_{-}, k_{-}, \omega) \exp[i(k_{-}x + k_{-}y + \omega t)] dk_{-} dk_{-} d\omega$

$$V(x, y, t) = \int f(\kappa_x, \kappa_y, \omega) \exp[i(\kappa_x x + \kappa_y y + \omega t)] d\kappa_x d\kappa_y dk_y$$

• Power spectrum $P(k_x, k_y, \omega) = ff^*$

 $k_h = (k_x^2 + k_y^2)$

October 13, 2005

Radial symmetric power spectrum

$$P(k_h,\omega) = \frac{1}{2\pi} \int_0^{2\pi} P(k_h \cos\phi, k_h \sin\phi, \omega)$$

Center for Solar-Terrestrial Research











