## Physics 681: Solar Physics and Instrumentation – Lecture 21

## **Carsten Denker**

**NJIT** Physics Department Center for Solar–Terrestrial Research



| Fig 1dS and Conducting Matter   |
|---|
| □ Magnetic features: sunspots, sunspot fine structure, filaments,<br>prominences, ... |
| □ Magnetic from the Sun occurs on all scales! (small flux tubes to mean field)        |
| □ Maxwell's equations (magnetic permeability μ = 4π × 10 <sup>-7</sup> Vs/Am)         |
| ∇ · B = 0   |
| ∇ × B = μj  |
| ∇ × E = - $\frac{dB}{dt}$   |
| □ Ohm's law   |
| Ü = σE with J = j and E = E + v × B   |
| ⇒ j = σ (E + v × B)   |
| □ Center for Solar-Terrestrial Research   |

\n- □ Induction equation\n 
$$
\frac{dB}{dt} = \nabla \times (\mathbf{v} \times \mathbf{B}) - \nabla \times (\eta \nabla \times \mathbf{B})
$$
\n
\n- □ Magnetic diffusivity\n 
$$
\eta = \frac{1}{\mu \sigma}
$$
\n
\n- □ Magnetic Reynolds number\n 
$$
R_m = \frac{vl}{\eta} = \frac{\tau_D}{\tau_A} \text{ with } \tau_D = \frac{l^2}{\eta} \text{ and } \tau_A = \frac{l}{\nu}
$$
\n
\n- □ High conductivity:\n 
$$
R_m \gg 1 \text{ or } \tau_D \gg \tau_A
$$
\n
\n- □ Electrical conductivity of a fully ionized gas\n 
$$
\sigma = \frac{32\epsilon_0^2 \sqrt{\pi} (2kT)^{3/2}}{\sqrt{m_e e^2 Z \ln \Lambda}} \gamma_E \text{ with } \Lambda = \frac{r_D}{p_0}
$$
\n
\n- November 15, 2005\n Center for Solar-Terrestrial Research

\n
\n

- $\Box$  In Λ ≈ 5 for the solar interior, In Λ ≈ 10 for the chromosphere, and ln Λ ≈ 20 for the corona
- **Conductivity is large:**

 $\sigma \approx 0.003 T^{3/2}$  A/Vm

- $\Box$  The temperature in the solar photosphere/chromosphere is low  $\Leftrightarrow$ hydrogen and helium are neutral BUT heavier elements with low ionization potentials still donate electrons  $\Rightarrow$  collisions of electrons with neutrals  $\Rightarrow$  conductivity is greatly reduced
- Conductivity for weak ionization

$$
\sigma = \frac{3e^2}{8S\left(2\pi m_{\rm e}kT\right)^{1/2}}\frac{n_{\rm e}}{n_{\rm n}}
$$

- □ Large horizontal gradients (e.g., sunspots)
- $\Box$  Collisions are rare and magnetic fields are strong  $\Leftrightarrow$  anisotropy
- □ Plasma turbulence reduces conductivity

November 15, 2005 Center for Solar-Terrestrial Research



| Example 1   | Example 2 |
|---|-----------|
| □ High conductivity $σ = 1$ V/A (in sunspots), typical velocity $v = 1$ km/s, and scale height $l = 100$ km ⇒ frozen field lines  |           |
| □ Total magnetic flux   |           |
| □ Differential flux at two instances separated by an infinitesimal time interval $dt$   |           |
| □ Differential flux at two instances separated by an infinitesimal time interval $dt$   |           |
| □ $\oint v - \Phi = dt \left[ \int \frac{dB}{dt} df + \int \frac{dB}{s} \cdot (v \times ds) \right]$  |           |
| with $B \cdot (v \times ds) = (B \times v) ds$ and $\int \frac{B \times v}{s} ds = -\int \left[ \nabla \times (v \times B) \right] \cdot df$  |           |
| □ $\frac{d\Phi}{dt} = \int \left[ \frac{dB}{dt} - \nabla \times (v \times B) \right] \cdot df$  |           |
| □ $\frac{d\Phi}{dt} = \int \left[ \frac{dB}{dt} - \nabla \times (v \times B) \right] \cdot df$  |           |
| □ $\frac{d\Phi}{dt} = \int \left[ \nabla \times (v \times B) \cdot \frac{d\Phi}{dt} \right] \cdot df$   |           |
| □ $\frac{d\Phi}{dt} = \int \left[ \nabla \times (v \times B) \cdot \frac{d\Phi}{dt} \cdot \frac{d\Phi}{dt$ |           |